

LASVET
Patent and Trademark Agency
Established 1992

28 September 2001

Patent Examiner

YOUR REF:

United States Patent & Trademark Office

OUR REF: 0968/P/JO

RE: TRANSLATION OF RUSSIAN SPECIFICATION

I, Jaak G. OSTRAT, am a qualified translator, well-versed in English and Russian.

I confirm that the Russian specification filed herewith, and identified below, was translated into English by me. By signing below, I also hereby declare that the specification filed herewith is a valid and accurate translation of the original Russian Specification.

1) U.S. Patent Application Number 09/885,645 filed on June 20, 2001

I, as an agent for the assignor, also confirm all inventors whose names appear on the Combined Declaration and Power of Attorney, filed herewith, signed the document fully aware of the words contained therein.

By signing below, I hereby verify that all statements made herein are correct and accurate to the best of my knowledge.



Signature of Translator

Jaak G. OSTRAT

Printed Name

28 September 2001

Date of Execution

W5
A1

METHOD OF TREATMENT OF MALIGNANT NEOPLASMS AND COMPLEX PREPARATION HAVING ANTINEOPLASTIC ACTIVITY

5

The invention relates to the field of medicine, oncology in particular, and can be used for chemotherapeutical treatment of oncological patients, suffering from different kinds of malignant neoplasms.

10

Methods of treatment of cancer by means of chemical preparations are known. According to mechanism of action chemical preparations are subdivided into alkylating agents, antimetabolites, alkaloids, antibiotics, hormones, immunomodulators and some others. In the course of treatment by different chemical preparations, the advantages and disadvantages thereof have been revealed.

Owing to comparatively small working concentrations, the chemical preparations affecting DNA are widely used. Alkylation of the vital DNA molecule leads to incapability of normal fission of cells and to their subsequent elimination. However, at the same time, DNA of normal non-cancerous cells is also subjected to alkylation. In order to reduce toxicity in relation to normal cells, when such preparations are applied, the methods for targeted delivery of preparations to cancerous cells by means of specific ligands, for example alpha-fetoprotein (AFP), are used. AFP is a transport protein, capable of delivering the substance bounded thereto to the cells having corresponding receptors. Such receptors abound on actively proliferating cancerous cells.

The method of treatment of primary liver cancer by means of intra-arterial injection of AFP in an amount of 2-20 mg, mainly 15 mg, five times at an interval of 7-12 days is known (Patent 35 RU No. 2,058,153, cl. A 61 K 38/00, published April 20, 1996, Bulletin No. 11).

The known method foresees the use of large amounts of AFP and distinguishes by long duration and limited extent of use, since the conditions for selecting patients for treatment by such a method are absence of jaundice and ascites, unoperability, absence of serious concurrent diseases.

The method of treatment of malignant neoplasms by means of intravenous injection of a complex preparation, comprising AFP and estrone-doxorubicin conjugate (Patent RU No. 2,026,688, cl. A 61 K 38/00, published January 20, 1995, Bulletin No. 2), is known. For preparation of the complex preparation impure AFP, an abortive material is used which is concentrated after rough purification to the content of AFP of 100 µg/ml and is then sterilized. Thereafter an antineoplastic antibiotic - doxorubicin is subjected to conjugation with a ligand - estrone in an equimolar ratio. Immediately 1-2 hours before intravenous injection the concentrate of AFP in an amount of 100 ml is mixed with 20 ml of the conjugate and thus one dose of the complex preparation for one injection (a single dose of AFP comprises 10 mg) is obtained. The overall course of treatment comprises 6 injections (three times every other day at an interval of a week).

The disadvantage of the known method is the use of high doses of AFP, the disadvantage of the known complex preparation is a labour-intensive procedure of obtaining the triple complex AFP-estrone-doxorubicin and impossibility of storage of the preparation for more than 2 hours.

30

The closest method to the present method - the prototype - is a method of treatment of primary liver cancer, comprising injection of the preparation doxorubicin dissolved in Lipiodol ultrafluid into the liver artery, while as the preparation doxorubicin doxorubicin-estrone dissolved in 96% ethyl alcohol at 70-76°C in a dose of 20-60 mg in 10-15 ml of Lipiodol

ultrafluid is used. 20 minutes prior to that a dose of 2-10 mg of AFP in 12-15 ml of physiological salt is injected into the liver artery; the repeated treatment is carried out after 3-4 weeks (Patent RU No. 2,065,307, cl. A 61 K 38/17, published 5 August 20, 1996, Bulletin No. 23).

The disadvantages of the known method are:

- labour-intensity of the method, connected with separated 10 injection of AFP and of doxorubicin-estrone complex;
- the use of high concentrations of the chemical preparations (AFP - 2-10 mg, doxorubicin-estrone complex - 20-60 mg), which can lead to toxic side reactions;
- high cost of treatment.

15 The closest complex preparation to the present complex preparation - the prototype - is a kit used for treatment of primary liver cancer and comprising doxorubicin-estrone in an amount of 20-60 mg in 15 ml vials, Lipiodol ultrafluid in two 20 ampules by 10 ml, AFP in an amount of 2-10 mg in 10 ml vials, physiological salt in volume of 15 ml in an ampule and 96% ethyl alcohol in volume of 5 ml in an ampule (Patent No. 2,065,307, cl. A 61 K 38/17, published August 20, 1996, Bulletin No. 23).

25 The disadvantages of the known kit are:

- labour-intensity and duration of obtaining the sterile doxorubicin-estrone complex as well as labour-intensity of preparation of the working solution of the given preparation: 30 the preparation is dissolved in 0.5-1.5 ml of 96% ethyl alcohol, heating to 70-76°C, the resultant solution is transferred into 10-15 ml of previously upwarmed Lipiodol ultrafluid, the suspension thus obtained is cooled to 32-37°C and is then injected into the liver artery under X-ray 35 television control;
- high concentrations of the preparations used in the kit;

- polycomponent nature of the kit.

The technical object of the group of the inventions is to simplify the known method, to lower the doses of the preparations to be injected and to reduce the costs of treatment.

The established object has been achieved by means of hereby proposed method of treatment of malignant neoplasms and a complex preparation having neoplastic activity for use in such treatment.

The realization of the method of treatment.

Depending on the character and seriousness of the disease, to a patient, together with the basic infusion-detoxication therapy, the complex preparation, comprising AFP in an amount of 0.07-0.15 mg, a polyene antibiotic in an amount of 4.2-7.0 mg and a pharmaceutically suitable filler, is parenterally injected once within twenty-four hours at an interval of three days in a course of 10 infusions.

The complex preparation having antineoplastic activity comprises the following components, in mg:

AFP	0.07 - 0.15
a polyene antibiotic	4.2 - 7.0
a filler	3.5 - 5.0

As a polyene antibiotic amphotericin B or nystatin is mainly used.

As a filler sugars are mainly used, for example glucose or synthetic polymers, selected from the group of polysaccharides, for example polyglykine, rheopolyglykine and dextran.

35

The complex preparation is obtained as follows: AFP with purity

not less than 98% is dissolved in distilled water or physiological salt in an amount of 0.07-0.15 mg/ml, whereto a polyene antibiotic in an amount of 4.2-7.0 mg/ml and then a filler in an amount of 3.5-5.0 mg/ml are added, after that the components are carefully mixed and the resultant mixture is then left to stand at 18-25°C for 10-12 hours. Thereafter the solution is sterilized by filtration, pre-packed in ampules or vials having a capacity of 1.2 or 3 ml and freeze-dried. The mass ratio of AFP, a polyene antibiotic and a filler is 1 : (60-100) : (50-70).

The complex preparation (conventional designation Reducin) is a powder of yellow colour, soluble in water, in physiological salt, in glucose solution and in other diluents (carriers), suitable for intravenous injection. One ampule of the complex preparation comprises: 0.07-0.15 mg of AFP, 4.2-7.0 mg of polyene antibiotic and 3.5-5.0 mg of corresponding filler. For preparation of the working solution the content of the ampule is dissolved in 2-3 ml of sterile water and is then transferred into 200 ml vial together with a carrier, suitable for intravenous injection.

The present method comprises injection to a patient of a new complex chemical preparation having antineoplastic activity and consisting of a vector part (AFP), specific to cancerous cells, and a nonspecific part, comprising a cytotoxic substance. As the latter an essentially new channel-forming and surface-active agent (SAA), namely a polyene antibiotic, for example amphotericin B or nystatin, which have not been previously used as antineoplastic remedies, is used. The targets for the new complex preparation are membranes of intracellular substructures. Such substructures include mitochondria, lysosomes, endoplasmatic reticula (EPR), nuclei, etc. In case of disorders in the function of membranes of substructures, a normal cell function, ensurable by the compartmentalization of different functions, is impossible. As

a result autodigestion of cancerous cells takes place according to the mechanism of induced apoptosis.

The use of AFP as a vector for targeted delivery of cytotoxic preparations to cancerous cells is known (see for instance patent RU No. 2,071,351, cl. A61 K 38/17, published January 10, 1997). In all known cases AFP is bounded with a cytotoxic part of the preparation by a chemical covalent bond, whereas in the present complex preparation AFP and SAA form a noncovalent complex, ensuring simultaneously the stability of the macromolecule during transportation and its functional independence in the process of cytotoxic action of the polyene antibiotic. The activity of the present complex preparation is based on the initiation of physiological reduction of cancerous cells according to the mechanism similar to that of apoptosis. As a rule, the natural autodestruction of the cells of tumor under the influence of the present complex preparation does not lead to intoxication of the organism and the effect of the treatment appears quickly.

The essential distinctive features of the present method, as compared with those of the prototype, are:

- to a patient the complex preparation, comprising AFP in an amount of 0.07-0.15 mg, a polyene antibiotic in an amount of 4.2-7.0 mg and a pharmaceutically suitable filler, is injected parenterally and simultaneously in a course of 10 injections (infusions) once in three days, which enables to simplify the known method, to lower the doses of the preparations to be injected and to reduce the costs of treatment.

The essential distinctive features of the present complex preparation, as compared with those of the known preparation, are:

- the complex preparation comprises additionally a surface-

active agent, namely a polyene antibiotic, mainly amphotericin B or nystatin, in an amount of 4.2-7.0 mg, which provides the complex with a new type of bond with a vector part - with a noncovalent bond, and a new mechanism of interaction with cancerous cells - with the membranes of the substructures of cancerous cells (membranes of lysosomes, EPR, mitochondria, nuclei), which, in its turn, improves the efficiency of the treatment and reduces side complications;

- the complex preparation comprises the components in optimal, experimentally selected amounts and ratios, namely: AFP - 0.07-0.15 mg, a polyene antibiotic - 4.2-7.0 mg, a filler - 3.5-5.0 mg, which enables to improve the efficiency of the treatment and substantially lower the doses of the active components used.

Being supported by the fact that no analogous method of treatment of malignant neoplasms and no analogous complex preparation having antineoplastic activity have been revealed, it may be concluded that the present group of inventions meet the requirements for patentability in respect of "novelty" and "inventive step".

The present method has been tested on 8 patients having IV clinical stage of oncological diseases, progressing after traditional ways of treatment. A full or a partial clinical effect has been achieved on 6 patients out of 8 (75%). The terms of remission were from 6 months to 1.5 years. In majority cases, for achieving a well-defined clinical effect it was sufficient to conduct one course of treatment.

The inventions are characterized by the following Examples.

Example 1

For preparation of the complex preparation Reducin 700 mg of AFP, 42 g of amphotericin B and 50 g of rheopolyglykine are dissolved in 1 litre of distilled water by mixing, subsequently

the volume is increased up to 10 litres. The resultant mixture is incubated at room temperature for 10-12 hours and is then subjected to sterilization by filtration through a membrane filter, pre-packed in 10,000 ampules or vials by 1 ml (a single dose) and thereafter freeze-dried in aseptic conditions. One ampule (vial) comprises 0.07 mg of AFP, 4.2 mg of amphotericin B and 5.0 mg of rheopolyglykine.

Example 2

10

The complex preparation Reducin is obtained analogously to Example 1, with the exception that to water solution 1 g of AFP, 50 g of amphotericin B and 40 g of polyglykine are added. As a result the preparation, comprising 0.1 mg of AFP, 5.0 mg of amphotericin B and 4.0 mg of polyglykine in a single dose, is obtained.

Example 3

20 The complex preparation is obtained analogously to Example 1, with the exception that to water solution 1.5 g of AFP, 70 g of amphotericin B and 30 g of dextran (molecular mass 100 kDa, Serva) are added. As a result the preparation, comprising 0.15 mg of AFP, 7.0 mg of amphotericin B and 3.0 mg of dextran in a single dose, is obtained.

Example 4

The complex preparation is obtained analogously to Example 1, 30 with the exception that to water solution 750 mg of AFP, 60 g of nystatin and 50 g of glucose are added. As a result the preparation, comprising 0.075 mg of AFP, 6.0 mg of nystatin and 5.0 mg of glucose in a single dose, is obtained.

35 Example 5

WILLIAM L. BROWN

A course of polychemotherapy was conducted according to the scheme CAF: On the 1st and 8th days intravenously 1 g of cyclophosphane, on the 1st and 8th days intravenously 500 mg of 5-fluorouracil, on the 2nd and 9th days of the treatment intravenously 40 mg of adriablastin. The treatment was accompanied by high toxicity without an expressed clinical effect.

25 Thereafter the patient was treated according to the present method. The complex preparation Reducin, comprising 0.07 mg of AFP, 4.5 mg of amphotericin B and 5.0 mg of glucose, was injected by infusion in a course of 10 injections once in three days. The condition of the patient has improved and became
30 satisfactory a week after the course of treatment had been finished. On the basis of the data of the radiology inspection of the organs of thorax favourable changes were registered, characterized by the decreased number of metastases in the lungs and the weakening of intensity of the shadows of dissemination.
35 The liquid in the pleural cavity was not inspected. Dyspnea on ascent has diminished and cough disappeared completely. During

Case No.	Plaintiff	Defendant	Amount	Verdict	Jury	Remarks
1	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
2	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
3	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
4	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
5	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
6	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
7	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
8	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
9	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
10	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
11	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
12	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
13	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
14	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
15	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
16	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
17	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
18	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
19	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
20	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
21	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
22	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
23	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
24	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
25	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
26	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
27	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
28	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
29	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
30	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
31	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
32	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
33	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
34	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
35	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
36	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
37	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
38	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
39	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
40	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
41	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
42	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
43	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
44	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
45	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
46	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
47	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
48	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
49	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
50	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
51	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
52	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
53	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
54	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
55	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
56	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
57	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
58	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
59	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
60	John Doe	Jane Smith	\$100.00	\$100.00	Yes	
61</						

5

15 The treatment program included a course of polychemotherapy according to the scheme CAMF. To the patient were injected 1 g of cyclophosphane on the 1st and 8th days, adriamicin on the 1st and 8th days, 50 mg of methotrexate on the 1st and 8th days, 5-fluorouracil on the 2nd and 9th days. The treatment was
20 accompanied by endotoxiosis and an abrupt worsening of the condition of the patient. No positive effects were detected.

6 days after the beginning of the treatment according to the
30 present method the diminishing of the sizes of subcutaneous
metastases were noted. By the end of the treatment rapid
improvement of the main disease was noted, which became apparent
from the twofold reduction of the size of all surface metastatic
nodes, the disappearing of pains in the lower jaw, the decreasing
35 of the rate of exudation in the right pleural cavity. Thrice
conducted right-sided pleural puncture proved the decreasing of

volume of exudate: before the treatment the volume of exudate was 600 ml, a week after the treatment - 350 ml, three weeks after the treatment - 20 ml. The improvement of function of central nervous system became apparent from restoration of normal swallowing function, restoration of gripping function of the left hand, clinically corresponding to the reduction of metastasis in the right hemisphere of brain.

The use of the present method of treatment of malignant neoplasms and the complex preparation having antineoplastic activity, as compared with the known method of treatment and the kit for use in such treatment, enables:

- to simplify the method of treatment by means of simultaneous parenteral injection of the complex preparation, comprising AFP and cytotoxic SAA in optimal ratios;
- to lower the doses of the used components: AFP 13-140 times, the cytotoxic component 3-14 times;
- to improve the efficiency of treatment by means of using the complex preparation having high specificity to growing cancerous cells and optimal composition of the complex preparation having a long term of storage (two years);
- to reduce the costs of treatment by means of using a small number of components in the complex preparation and low doses of the chemical preparations to be used.

Thus, while using the present method of treatment of malignant neoplasms two biological mechanisms are realized. The first of them comprises the targeted delivery of the cytotoxic agent by means of AFP to tumorous cells. The second one comprises the directed destruction of the tumorous cells because of destruction of the intracellular structures, in particular EPR and lysosomes. This may be accompanied by autodigestion of tumorous cells, caused by the enzymes of hydrolysis, comprising in the lysosomes, according to the mechanism of targetably induced apoptosis. The directed reduction affects much less the cells of blood-forming,

30

Claims:

sub
A2
1. A method of treatment of malignant neoplasms, comprising
5 injection of alpha-fetoprotein (AFP), characterised in that AFP
is injected as a component of the complex preparation comprising
additionally a polyene antibiotic and a filler in ratio of 1:
(60-100) : (50-70), and the preparation is injected parenterally
in a course of 10 injections once in three days.

10

2. A complex preparation having antineoplastic activity,
comprising AFP, a cytotoxic substance and a filler, characterized
in that the preparation comprises a polyene antibiotic as a
cytotoxic substance and polysaccharide or sugar as a filler in
15 the following quantitative ratio of the components, in mg:

AFP	0.07 - 0.15
a polyene antibiotic	4.2 - 7.0
a filler	3.5 - 5.0

20 3. A complex preparation of claim 2, characterized in that the
preparation comprises amphotericin B or nystatin as a polyene
antibiotic.

4. A complex preparation of claim 2, characterized in that the
25 preparation comprises polysaccharide, selected from the group
consisting of polyglykine, rheopolyglykine and dextran, or sugar,
mainly glucose as a filler.

30

add
A3

add
B2

09/885645

Abstract

Use: medicine, oncology in particular.

5

Technical object: simplification of the known method, lowering of the doses of injectable medicines and reducing the costs of treatment.

10 Essence of the invention:

Depending on the character and seriousness of the disease, to a patient, together with the basic infusion-detoxication therapy, the complex preparation, comprising AFP in an amount of 0.07-0.15 mg, a polyene antibiotic, mainly amphotericin B or nystatin, in an amount of 4.2-7.0 mg and a filler, is parenterally injected once within twenty-four hours at an interval of three days in a course of 10 infusions.

20 The complex preparation having antineoplastic activity comprises the following components, in mg:

AFP	0.07-0.15
a polyene antibiotic	4.2-7.0
a filler	3.5-5.0

25

As a filler polysaccharide from the group of rheopolyglykine, polyglykine, or sugar, e.g. glucose is mainly used.

Positive effect: Simplification of the method, lowering of the doses of injectable medicines and reducing the costs of treatment. The complex preparation distinguishes by high efficiency of antineoplastic activity, a small number of components, simplicity of preparation and a long term of storage.